

Submitted to ApJ June 7, accepted March 17, 2007

New Photometry and Spectra of AB Doradus C: An Accurate Mass Determination of a Young Low-Mass Object with Theoretical Evolutionary Tracks¹

Laird M. Close², Niranjana Thatte³, Eric L. Nielsen², Roberto Abuter⁴, Fraser Clarke³, & Matthias Tecza³

lclose@as.arizona.edu

ABSTRACT

We present new photometric and spectroscopic measurements for the unique, young, low-mass evolutionary track calibrator AB Dor C. While the new Ks photometry is similar to that previously published in Close et al. (2005) the spectral type is found to be earlier. Based on new H & K IFS spectra of AB Dor C (Thatte et al. 2007; paper I) we adopt a spectral type of $M5.5 \pm 1.0$ for AB Dor C. This is considerably earlier than the $M8 \pm 1$ estimated in Close et al. (2005) and Nielsen et al. (2005) yet is consistent with the $M6 \pm 1$ independently derived by Luhman & Potter (2005). However, the spectrum presented in paper 1 and analyzed here is a significant improvement over any previous spectrum of AB Dor C. We also present new astrometry for the system which further supports a $0.090 \pm 0.005 M_{\odot}$ mass for the system. Once armed with an accurate spectrum and Ks flux we find $L = 0.0021 \pm 0.0005 L_{\odot}$ and $T_{eff} = 2925^{+170}_{-145} \text{K}$ for AB Dor C. These values are consistent with a $\sim 75 \text{ Myr } 0.090 \pm 0.005 M_{\odot}$ object like AB Dor C according to the DUSTY evolutionary tracks (Chabrier et al. 2000). Hence masses can be estimated from the HR diagram with the DUSTY tracks for young low-mass objects like AB Dor C. However, we cautiously note that underestimates of the mass from the tracks can occur if one lacks a proper (continuum preserved) spectra or is relying on NIR fluxes alone.

⁰ Based on observations made with ESO Telescopes at the Paranal Observatories under programme ID 276.C-5013.

²Steward Observatory, University of Arizona, Tucson, AZ 85721

³Dept. of Astrophysics, DWB, University of Oxford, Keble Road, Oxford OX1 3RH, U.K.

⁴European Southern Observatory, Garching, Germany

Subject headings: instrumentation: adaptive optics — binaries: general — stars: evolution — stars: formation — stars: individual (AB Doradus C) — Brown Dwarfs — extrasolar planets

1. Introduction

There is currently great interest in the direct detection of extra-solar planets, brown dwarfs, and very low-mass stars. The study of such young, low-mass objects has been yielding increasingly fruitful science, yet the field remains dependent on evolutionary models to properly interpret the data that are collected from these objects. In particular, mass, while a fundamental property, is very rarely measured directly (through orbital dynamics), and instead must be inferred from theoretical tracks (e.g., Burrows, Sudarsky, and Lunine 2003, Chabrier et al. 2000). It is thus of great interest to discover calibrating objects that can link a dynamically measured mass with observables such as accurate NIR (1-2.5 μm) fluxes and spectral types.

AB Dor A was suspected of having a low-mass companion due to VLBI and Hipparcos measurements of an astrometric wobble (Guirado et al. 1997). Recently Close et al. (2005), reported the first direct detection of this low-mass companion (AB Dor C) to the young star AB Dor A, along with the first measurements of the JHKs fluxes, spectral type, and dynamically determined mass of AB Dor C. Upon comparing these results with the predictions of Chabrier et al. (2000), they found the models to be systematically over-predicting the fluxes and temperature of AB Dor C (especially at J & H), given a system age of 50 Myr. Put another way, the model masses seem to be underestimating the mass of a low-mass object given its age, J & H NIR fluxes, and spectral type. Since the publication of these results, another calibrating object has been reported by Reiners, Basri, and Mohanty (2005): USco CTIO 5. While this equal-mass binary is younger (~ 8 Myr) and more massive (total mass $\geq 0.64 M_{\odot}$) than AB Dor C, Reiners et al. find the same trend of models under-predicting masses based simply on photometric and spectroscopic data applied to the HR diagram. A similar trend for such masses was previously noted by Hillenbrand and White (2004). Moreover, this trend has been theoretically predicted for higher masses by Mohanty, Jayawardhana, and Basri (2004), and by Marley et al. (2005) for planetary masses. Hence it is critically important to accurately calibrate the evolutionary tracks to determine if there are systematic errors. In particular, obtaining an accurate T_{eff} from spectra of low-mass, young, objects is challenging and makes comparison to evolutionary tracks difficult (Chabrier et al. 2005).

Recently Nielsen et al. (2005) have obtained new orbital epochs and confirmed the previous mass of AB Dor C of $0.090 \pm 0.005 M_{\odot}$ with the technique of Guirado et al. (2006).

Hence AB Dor C has a uniquely well known dynamical mass for a young (~ 75 Myr) low-mass object. Only GL 569Ba/Bb has a similarly well determined mass but is thought to be somewhat older (age ~ 300 Myr; Zapatero Osorio et al. 2005), though a younger age (~ 100 Myr) and binarity of GL 569 Ba has recently been proposed (Simon et al. 2006). Also the lack of lithium in the spectrum of GL 569B (Zapatero Osorio et al. 2006) is surprising. Hence, there is a need to quantify the AB Dor A/C system as closely as possible to have an additional calibrator for low-mass young evolutionary models.

The age of AB Dor (and its associated moving group) is also somewhat uncertain, Luhman et al. (2005) argue for an older age of the AB Dor group of 75-150 Myr while Nielsen et al. (2005) find 75 ± 25 Myr. Recently Lopez-Santiago et al. (2006) argue for a 50 Myr age for the core of the AB Dor moving group (which includes AB Dor A) as did Zuckerman et al. (2004) in the original “AB Dor moving group” paper. Recently Janson et al. (2007) find an age range of 50-100 Myr. Here we adopt an “average” age of 75 Myr for the system.

There is also uncertainty in the spectral type of AB Dor C. The small separation between A & C of only $0.155''$ combined with the > 100 contrast made an accurate spectral type difficult to measure in the dataset of Close et al. (2005). A reanalysis of the Close et al. spectra by Nielsen et al. (2005) suggested a spectral type of $M8 \pm 1$. However, an independent re-analysis of these data by Luhman & Potter (2005) suggested a spectral type of $M6 \pm 1$. The importance of this temperature is paramount to plotting AB Dor C on the HR diagram to calibrate the accuracy of the evolutionary tracks. These past papers highlight the difficulty in trying to determine the spectral type of a faint companion within $0.16''$ of a bright star with an AO-fed long-slit spectrograph. Indeed none of these past reductions of the AO long-slit spectra were able to preserve the continuum of AB Dor C. Hence the true spectrum of C could not be accurately determined until now.

In the companion paper to this one (Thatte et al. 2007; hereafter Paper I) we report on new, excellent integral field spectroscopy (IFS) of AB Dor A & C, observed with the SINFONI instrument at the VLT. Using the new data analysis technique of PSF scaling and differencing (PSD), we were able to effectively eliminate all contamination from AB Dor A, to produce a very high signal-to-noise spectrum of AB Dor C, which also preserves the continuum. The PSD technique is able to achieve very high contrast (~ 9 mags at $0''.2$) without a coronagraph, and without any prior assumptions about the spectral characteristics of the companion object (see Paper 1 for more information about the IFS reduction, and the PSD technique).

In this paper we present new VLT science archive Ks photometry and astrometry for AB Dor C and a more accurate H & K spectrum and a new temperature of AB Dor C.

Then we compare the accuracy of the mass of AB Dor C predicted by the popular DUSTY theoretical tracks (Chabrier et al. 2000).

2. OBSERVATIONS & REDUCTIONS

The AB Dor system (A, C, and their companion BaBb – 8.87'' distant) was observed with the VLT NACO AO system (Lenzen et al. 2003; Rousset et al. 2002) on Jan 7, 2005 by R. Neuhauser et al. in the Ks band. We have reduced these archive data with our NIR AO data reduction IRAF pipeline (Close et al. 2002, 2003).

The Ks images were fully flat-fielded, sky & dark subtracted, bad pixel cleaned, and aligned and medianed (see Close et al. 2003 for more details about our AO pipeline). Only the first 18 200x0.347 second Ks images were reduced since the seeing became worse after these first 18 images.

We also observed AB Dor A & C with the SINFONI IFS as described in paper I. See paper I for extensive details of the IFS observations & reductions.

3. ANALYSIS

3.1. The Ks Flux and Astrometry of AB Dor C

To accurately measure the flux of a tight faint companion is never trivial. However, by January 2005 AB Dor C had moved out to a 0.22'' separation from AB Dor A. While this may seem a very small separation it is considerably better than the 0.155'' separation during the Feb 2004 discovery images of Close et al. (2005). Hence, we should be able to better gauge the flux of AB Dor C at this latter epoch.

To measure the brightness of AB Dor C we utilized the unsaturated PSF image (in the narrow band 2.12 micron filter) of AB Dor A which was taken just before the AB Dor Ks data set. This 2.12 μm “PSF” image was taken at the same airmass and seeing as the Ks images. Also, the exposure times were the same (DIT=0.347 seconds, NDIT=200 coadds) for the PSF and Ks images. The FWHM of the PSF appeared similar to that of AB Dor C. So we have some confidence that this was a good PSF.

We shifted (with a cubic spline) and scaled this PSF image until subtracting it led to the flux from C being completely removed from the reduced Ks images. However, as is clear from Figure 1, there is some residual flux or “super-speckles” surrounding C’s position from

A. Therefore, it is impossible to absolutely determine the flux of C due to some uncertainty in A’s PSF. We adopt the mean flux of AB Dor C as that where the residual flux of A at the position of C is equal to that 180 degrees on the other side of the A PSF (see bottom left image in figure 1). This is a reasonable assumption since the super speckles in the NACO PSF are mainly due to phase errors which transform to symmetric pairs on either side of the PSF. Here we have made the conservative assumption that the possible range of C’s flux should be from completely removing all residual flux at the position of C (Fig 1: lower right) to assuming there is a slightly brighter super-speckle at C’s position (Fig 1; upper right).

3.1.1. Photometry of AB Dor C

The Ks flux of AB Dor B is known from the 2MASS Catalog to be $Ks_B = 7.435$ (corrected for the 0.095 mag of contamination of B due to A in the 2MASS 4'' aperture photometry). Hence by calibrating the mean delta Ks of the PSF to BaBb (which varied by $< 2\%$ over a range of 1.5, 2.0, & 2.5'' apertures) we derive a $Ks = 9.50 \pm 0.16$ mag for C. This flux is derived assuming a symmetric PSF while the large errorbars are derived assuming minimal symmetry in the AB Dor A PSF and adding in the maximum errors of the aperture photometry (see the top right and bottom right images in Figure 1). Hence, the true flux of AB Dor C should certainly fall inside this range. Our new value of $Ks = 9.50 \pm 0.16$ is very similar to the $9.45^{+0.060}_{-0.075}$ mag measured by Close et al. (2005). Our conservative assumption of minimal PSF symmetry around AB Dor A leads to our photometric errors being larger than those of Close et al. who assumed the PSF was mainly symmetric. Our new photometry is also consistent (if a bit brighter and more precise) than the $9.79^{+0.25}_{-0.33}$ mag independently determined by Luhman & Potter (2005) from the Close et al.(2005) dataset. Moreover, we obtain a $Ks \sim 9.59$ mag from the IFS data-cube in paper 1 – adding further confidence to our Ks band flux of C. Hence, there is now reasonable agreement from three different epochs that $Ks = 9.5 \pm 0.16$ for AB Dor C.

From the Hipparcos distance of 14.9 ± 0.1 pc and the -2.82 ± 0.15 mag BC_K of Allen et al. (2003)(appropriate for M5.5) and noting that for an M5.5 $Ks - K \sim 0.03$ mag (Daemgen et al 2006) we derive a luminosity of $L = 0.0021 \pm 0.0005 L_\odot$ from the observed $Ks = 9.50 \pm 0.16$ mag of AB Dor C.

3.1.2. Astrometry of the AB Dor A/C system

From our PSF fitting AB Dor C is found to be $0.219 \pm 0.008''$ at $PA = 155.92 \pm 0.5^\circ$ in good agreement with the orbit of Nielsen et al. (2005) which predicts a separation of $0.2186''$ at $PA = 155.175^\circ$ for Jan 7, 2005 (2005.0170). This good agreement gives us further confidence in the orbital solution of Nielsen et al. (2005). Moreover, we have another (even later) Jan 24 2006 (2006.0660) epoch of observation from the IFS dataset of Paper I (see Table 1). This additional data-point also fits the orbital solution of Nielsen et al. (2005) very well. Figures 2 & 3 illustrate the quality of the orbital solution of Nielsen et al. (2005). Note that this orbit was mainly determined by the reflex motion of A from ~ 1 mas Hipparcos/VLBI astrometry of Guirado et al. (2006) (for clarity these Hipparcos/VLBI data-points are not shown in Figure 2).

Recently Jeffers, Donati & Cameron (2007) have estimated the radial velocity variations of the AB Dor A/C system from 1988 to 1994. They find a reasonable fit of our older Close et al. (2005) orbital solution to their radial velocity measurements. The more recent orbital solution of Nielsen et al. (2005) should be similar. This further proof of the quality of the A/C orbital solution.

3.1.3. Astrometry of the Ba Bb system

We were also able to study the companion binary BaBb (a tight $0.07''$, $PA = 238.6 \pm 0.3$ binary discovered by Close et al. 2005) some $9''$ away is visible in the reduced image. On Jan 7, 2005 we find the BaBb binary is now $0.060 \pm 0.003''$ in separation at $PA = 246 \pm 2$ (with $\Delta K_s = 0.22$ mag), while Ba is $8.87 \pm 0.10''$ at $PA = 346.31 \pm 0.5$ w.r.t. A (see Table 1 for more details). The small change in position of Bb w.r.t. Ba & A since the Feb. 2, 2004 Close et al. (2005) observations definitively proves BaBb is a tight binary itself bound to A. Previous observations show that B itself is in orbit around A (Guirado et al. 2006).

Although we know relatively little about the orbit of Bb around Ba we know AB Dor Ba/Bb has a projected separation of 0.9 AU, and for an almost equal mass binary with combined spectral type $\sim M4$ (Martin & Brandner 1995; see Table 1), this is consistent with a Keplerian orbit with a period of order one year. We have also obtained an acquisition image from the ESO VLT archive between the two observational epochs discussed here. This image (from ESO program 073.C-0469(B), Chauvin et al.) was taken on September 24, 2004: about seven and a half months after the Close et al. 2005 dataset, and about three and a half months prior to the Neuhaser et al. dataset. In this image, taken with the same NACO system, using the same S13 camera with identical platescale, the same Ks filter, and AB

Dor Ba/Bb having a similar FWHM compared to future and past epochs, the appearance of the binary is drastically different. Unlike the Feb. 2004 and Jan. 2005 images, where the two components of the binary are clearly distinguishable, the Sep. 2004 image appears to be of a single star. This short-period variation, combined with the similar separations and position angles observed between Feb. 2004 and Jan. 2005 seem to constrain the orbit of the AB Dor Ba/Bb system to approximately 11 months. Future monitoring of this system is required to obtain more detailed orbital parameters.

3.2. The Spectral Type of AB Dor C

As is outlined in paper I an excellent $R\sim1500$ H and Ks spectrum of AB Dor C was obtained with the SINFONI IFS. Here we attempt to place our spectrum on a spectral sequence. Unfortunately there simply are no good H & K template spectra for late M stars of ~75 Myr of age. Hence, we have endeavored to fit our spectrum to young (~ 1 Myr; Gorlova et al. 2003) and old field M stars (Cushing et al. 2005). As is clear from figures 4 - 7 there is pretty good agreement that the spectrum is similar to M5 or a M6.

We feel the spectra presented here are superior to those previously published by Close et al. (2005); Nielsen et al. (2005); and Luhman & Potter (2006) for several reasons. All previous published spectra were different reductions of the long-slit NACO K spectra obtained by Close et al. (2005). These AO-fed long slit spectra suffered from many of the drawbacks of AO-fed long-slit spectra. First, the very small $0.085''$ slit used in the older spectra was only roughly aligned with the binary, due to flexure etc. Hence, there are different slit losses for each spectrum obtained as AB Dor was nodded along the slit. Moreover, as the core FWHM of the AO PSF decreases as λ increases, hence slit losses are also function of the wavelength as well as the centering error. Finally there may have been some error in the PA angle of the slit and so there may be different slit losses between AB Dor A and C. Also the very small separation of $0.15''$ between A & C made these spectra very difficult to reduce (in fact many of the individual spectra had to be dropped from the reduction since no manner of subtracting A could reveal a non-noise signal at the position of C). Indeed, Luhman & Potter (2006) and Nielsen et al. (2005) only use $\sim 50\%$ of the spectra obtained to try and detect a signal from C. In all these previous spectra the flux from AB Dor A swamped the light from C (just $0.15''$ away); this is not surprising given the $\sim 0.2''$ images obtained with the NACO spectrograph that night. In the end, none of the three previous efforts at reducing this Close et al. (2005) spectral dataset could obtain any meaningful K continuum of AB Dor C (and no H band spectra was attempted). So all these past efforts simply fit the "continuum" with a polynomial and attempted to remove

as much of it as possible (in Close et al. 2005 the continuum was restored with a standard). Therefore, no spectrum of AB Dor C with the original continuum intact have been published to date.

Without continua it is very hard to be sure that one is not either over or undersubtracting A from C. Hence, strengths of the CO and Na absorption lines are somewhat suspect and uncertain. This explains, in part, the significant dispersion in the spectral types determined from the three different reductions of the rather poor quality AO long slit spectra. To better understand what the real spectral type of AB Dor C is we have approached this problem with a new (more advanced) technique of PSD IFS reduction.

Our IFS spectrum in this paper is superior for several reasons. One is that C was a more favorable $0.20''$ away from A at the time of the IFS observation. Another is that use of the IFS allowed significantly better removal of the contaminating PSF wing of A (at every wavelength) of the IFS datacube by PSF subtraction (see paper 1 for full details). Moreover, we could easily check in each IFS "image" of how well we had removed the wing of A from the position of C (this is nearly impossible to do with long slit spectra). Also, by effectively integrating longer we were able to obtain $S/N \sim 40$. In addition, we have obtained H band as well as K band spectra. Increasing the spectral range through H helps determine a more accurate spectral type. Finally, use of the the IFS eliminates slit losses, and has allowed us to produce excellent spectra of AB Dor C in the H and K bands with a stable continuum across both bands. To highlight the improvement we directly compare our new spectra with that of Nielsen et al. (2005) in figure 2 of paper 1.

We believe that the points raised above make clear why our H & K IFS continuum-preserved spectrum is superior to previously published (non-continuum preserved) K spectra of AB Dor C. However, it is still difficult obtaining a perfect spectral type fit for AB Dor C since it has an age (and so surface gravity) different from that of published M standards. For example, in Fig 5 it is clear that M4 is not the correct spectral type since the Na lines of Gl 213 fit the AB Dor C spectra too well – and since AB Dor C is younger than Gl 213 it should have *weaker* Na I lines than a field dwarf of the same spectral type (see Gorlova et al. 2004 for example). In fact, the M5 field dwarf Gl 51 fits the CO lines the best (including an exceptionally good fit to the pseudo-continuum past 2.3 microns). Since CO is not strongly gravity dependent (Gorlova et al. 2004) there is evidence that the temperature of AB Dor C must be close to that of Gl 51 (M5).

Spectra presented in previous studies all likely had too much strength in the Na I line compared to CO (which is unlikely for a low surface gravity object). In this study it is now clear that in the K band data AB Dor C appears as a low surface gravity M5 (fitting the CO well but weaker in the Na I lines). Moreover, our K-band continuum follows that of Gl 51

(M5) or Gl 406 equally well, yet that of an M4 (too hot) or M7 (too cold) can be excluded at $> 1\sigma$ of our estimated noise level (see our 1σ errorbars in figures 5 & 6). Hence, there is excellent evidence from Fig 5 that the spectral type of C must be in the range of M5-M6.

Further evidence comes from Fig 6 (our H band spectra). Here we see the Mg & Al lines between 1.65-1.75 microns of the M3 & M4 dwarfs are not seen in the cooler AB Dor C atmosphere. Moreover, AB Dor C’s continuum shape is nicely matched by the M5 and M6 templates from 1.58-1.8 microns. Whereas the bluest part of H has the lowest Strehl and likely some slight contamination from the hotter A component – heating up our continuum slightly at the bluest part of H band. The M7 continuum appears too cool. Hence, we see in the H band the best fit is to a M5-M6 spectral type. Therefore, we determine that the gravity independent line strengths (like CO) and the continuum of AB Dor C are best fit by field M5-M6 dwarfs. In summary, we conservatively estimate a spectral type of $M5.5 \pm 1.0$ for AB Dor C.

Our new spectral type of $M5.5 \pm 1.0$ for AB Dor C is considerably earlier than the $M8 \pm 1$ measured in Close et al. (2005) and Nielsen et al. (2005) yet is consistent with the $M6 \pm 1$ independently derived by Luhman & Potter (2005). In summary, our continuum is preserved and the S/N (~ 40), wavelength range, and resolutions (~ 1500) are all much higher, hence the spectrum presented here is a significant improvement over any previous spectrum of AB Dor C.

Using the dwarf temperature scale of Leggett et al. (1996) we find a temperate range of $T_{eff} = 2925^{+170}_{-140}$ K for AB Dor C. However, this error does not take into account a possible ~ 150 K systematic error in this popular temperature scale. Hence the final total errors could be as high as 2925^{+226}_{-205} K.

4. Comparison to Models

In figure 7 we compare our observed values to those predicted by the cooling curves of the DUSTY models. Note how the J and H values appear to be fainter than the models would predict (assuming a 75 Myr age for AB Dor; Nielsen et al 2005), this result is similar to that of Close et al. (2005). Yet there is good agreement with the Ks (as was also the case in Close et al. 2005). Moreover, our higher T_{eff} is in better agreement with the cooling curves than in Close et al. (2005).

It is interesting to note that the J-Ks for AB Dor C is $\sim 1.26 \pm 0.21$ which is surprisingly red for M5.5 object (which are typically closer to $J - Ks \sim 1.0$ in the Pleiades). Hence, it is possible that the J band flux measured by Close et al. (10.76 ± 0.1) or Luhman & Potter

($10.72^{+0.40}_{-0.63}$) may be systematically too faint. We caution that it is problematic to obtain high-quality high-contrast images in the J band with AO and so, in general, weight should not be placed on J band fluxes of faint companions since lower Strehls lead to a higher chance of a poor flux calibration. In any case, the mean J flux of Close et al. (2005) would only have to be brightened by ~ 0.2 mag to give consistent colors with M5.5 Pleiades objects. This would lead to slightly better agreement with the models especially at the oldest ages suggested for AB Dor of 75-150 Myr by Luhman et al. (2005)).

As is mentioned in Luhman et al. (2005) a good technique for indirectly determining the mass of an object is to place the object in the HR diagram and from the evolutionary tracks estimate a mass. In figure 8 we show how the new spectral type of M5.5 is in good agreement with the DUSTY tracks in the HR diagram for a $0.090M_{\odot}$ object like AB Dor C. Hence, the combination of a careful measurement of the Ks flux and spectral type can allow the mass of a young low-mass object like AB Dor C to be estimated from the DUSTY tracks. However, underestimates of the mass can occur without proper (continuum preserved) spectra or relying on just H and J fluxes alone.

The points with error bars in Figure 8 mark all low-mass young objects with well determined dynamical masses. The thick diagonal lines represent the rough disagreement between the measured luminosity and temperature to the values actually predicted by the DUSTY models for the true mass of the object. One can see that for AB Dor C and the older (~ 300 Myr) Gl 569 B system (Zapatero Osorio et al. 2004), the offset between observation and the tracks are within the 1σ uncertainties (especially considering the uncertainty in the ages for these systems).

However, in the recently discovered eclipsing binary 2MASS J05352184-0546085 (Stassun et al. 2006) in the Orion Trapezium star formation region (~ 1 Myr age) the $0.034 \pm 0.0027M_{\odot}$ secondary is *hotter* than the $0.0541 \pm 0.0046M_{\odot}$ M6.5 primary. Hence, there is an increase in uncertainty in the models for very young (> 10 Myr) low-mass ($< 0.040M_{\odot}$) objects. However, the derived mass for 2M0535A would only be low by $\sim 25\%$ (just consistent with the 1 Myr track), whereas the derived mass for 2M0535B from the 1 Myr track would be about 200% too high (a similar trend was predicted by Mohanty et al. 2004).

We thank Mike Cushing and Nadja Gorlova for providing spectra of many young, low-mass objects. The Ks data was from observations made with the European Southern Observatory telescopes obtained from the ESO/ST-ECF Science Archive Facility under program ID 074.C-0084(B).

This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and

Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

LMC is supported by an NSF CAREER award and the NASA Origins of Solar Systems program. ELN is supported by a Michelson Fellowship. NT, MT and FC are supported by a Marie Curie Excellence Grant from the European Commission MEXT-CT-2003-002792 (SWIFT).

REFERENCES

- Allen, P.R., Trilling, D.E., Korner, D.W., Reid, I.N. 2003 ApJ 595, 1222.
- Baraffe, I., Chabrier, G., Allard, F., & Hauschildt, P.H. 1998 A&A 337 403.
- Burrows, A., Sudarsky, D., & Lunine, J.I. 2003 ApJ 569, 587.
- Chabrier, G., Baraffe, I., Allard, F., & Hauschildt, P.H. 2000 ApJ 542, 464.
- Chabrier, G., Baraffe, I., Allard, F., & Hauschildt, Invited review, "Resolved Stellar Populations", Cancun, april 2005. [astroph-0509798](#)
- Close L.M., Siegler, N., Potter, D.; Brandner, W., Liebert, J. 2002 ApJ 567, L53
- Close L.M., Siegler, N., Freed, M., Biller, B. 2003 ApJ 587 407
- Close, L.M., Lenzen, R., Guirado, J.C., Nielsen, E. L., Mamajek, E.E., Brandner, W., Hartung, M., Lidman, C., Biller, B. 2005 Nature 433, 286.
- Cushing, M.C., Rayner, J.T., & Vacca, W.D. 2005 ApJ 623, 1115.
- Gorlova, N.I., Meyer, M.R., Rieke, G.H., & Liebert, J. 2003 ApJ 593, 1074.
- Guirado, J.C., Reynolds, J.E., Lestrade, J.-F., et al. 1997 ApJ 490, 835G.
- Guirado, J.C., Marti-Vidal, I., Marcaide, J.M., Close, L.M., Algaba, J.C., Brandner, W., Lestrade, J.-F., Jauncey, D.L., Jones, D.L., Preston, R.A., and Reynolds, J.E. 2006 A&A 446 733.
- Hillenbrand, L.A. & White, R.J. 2004 ApJ 604, 741.
- Janson, M., Brandner, W., Lenzen, R., Close, L., Nielsen, E., Hartung, M., Henning, T., Bouy, H. 2007, A&A 462, 615

- Jeffers, S. V., Donati, J.-F., Collier Cameron, A. 2007 MNRAS in press Online Early 02/2007
- Leggett S.K., Allard, F., Berriman, G., Dahn, C.C., & Hauschildt, P.H. 1996, ApJS, 104, 117
- Lenzen, R. et al. 2003, SPIE 4841, 944.
- Lopez-Santiago, J. Montes, D., Crespo-Chacon, I. & Fernandez-Figueroa, M.J. 2006, ApJ 643, 1160
- Luhman, K.L. 1999 ApJ 525, 466.
- Luhman, K.L., Stauffer, J.R. & Mamajek, E.E. 2005 ApJ 628, 69.
- Luhman, K.L. & Potter D. 2006 ApJ 638 887
- Marley, M.S., Fortney, J., Hubickyj, O., Saumon, D., et al. 2005 Presentation, 2005 Winter Conference on Astrophysics.
- Martin, E.L. & Brandner, W. 1995, A&A 294 744
- Martin, E.L., Brandner, W., Bouvier, et al. 2000 ApJ 543, 299
- Martin, E.L., Rebolo, R. & Zapatero-Osorio, M.R. 1996 ApJ 469, 706
- Mohanty, S., Jayawardhana, R., & Basri, G. 2004 ApJ 609, 885
- Nielsen, E.L., Close, L.M., Guirado, J.C., Biller, B.A., Lenzen, R., Brandner, W., Hartung, M., and Lidman, C. 2005 AN 326 1033.
- Reiners, A., Basri, G., and Mohanty, S. 2005 ApJ 634 1346.
- Rousset, G. et al. 2002, SPIE 4389, 140.
- Simon M. et al. 2006 ApJ 644, 1183
- Stassun, K. G., Mathieu, R. D., Valenti, J. A. 2006, Nature, 440 311.
- Thatte N., Abuter, R., Tecza, M., Nielsen, E., Clarke, F.J., & Close, L.M. 2007 MNRAS in press (PAPER I), astro-ph/0703565
- Zapatero Osorio, M. R., Lane, B.F., Pavlenko, Ya., Martin, E.L., Britton, M., & Kulkarni, S.R. 2004 ApJ 615, 958
- Zapatero Osorio, M. R. et al. AN 326 948

Zuckerman, B., Song, I., & Bessell, M.S. 2004 ApJ 613L, 65

Table 1. The AB Dor System

Parameter	A	C	Ba	Bb
K_s magnitude	4.686 ± 0.016	9.50 ± 0.16	8.08 ± 0.20	8.30 ± 0.20
Separation ^a to A	–	$0.219 \pm 0.008''$	$8.87 \pm 0.10''$	–
PA ^a w.r.t A	–	$155.92 \pm 0.50^\circ$	$346.31 \pm 0.50^\circ$	–
Separation ^a to Ba	–	–	–	$0.060 \pm 0.003''$
PA ^a w.r.t. Ba	–	–	–	$246 \pm 2^\circ$
Separation ^f to A	–	$0.202 \pm 0.010''$	–	–
PA ^f w.r.t A	–	180.78°	–	–
Period w.r.t A (yr)	–	11.74 ± 0.07^c	$1400\text{--}4300^b$	–
Period w.r.t Ba (yr)	–	–	–	~ 0.9
Spectral Type	K1	$M5.5 \pm 1.0$	$M3.5 \pm 1.5$	$M4.5 \pm 1.5$
T_{eff} (K)	5081 ± 50	2925_{-145}^{+170}	3265 ± 245	3095 ± 255
Luminosity ($L_\odot$)	0.388 ± 0.008	0.0021 ± 0.0005	0.008 ± 0.002	0.006 ± 0.002
Mass ($M_\odot$)	0.865 ± 0.034	0.090 ± 0.005^c	$< 0.25^b$	$< 0.15^b$
System Age (Myr)	75 ± 25^d			
System Distance ^e (pc)	14.9 ± 0.1			

^aobservations made on Jan 7, 2005 UT (2005.0170)

^bOrbital calculations of Guirado et al. (2006)

^cOrbital solution of Nielsen et al. (2005)

^dAge solution of Nielsen et al. (2005)

^eHipparcos distance (15.06 ± 0.11 pc is derived by Hipparcos & VLBI in Guirado et al. (2006))

^fobservations made on Jan 24, 2006 UT (2006.0660)




Fig. 1.— Top left: AB Dor C and A (unsharp masked). Top right: After subtracting the PSF scaled by 4.5x ($K_s=9.66$ mag) there is too much residual light at the location of C – hence this is a lower limit to the flux of AB Dor C. Bottom left: A more optimal subtraction leaving the same amount of flux on either side of A ($K_s=9.50$ mag). Bottom right: An over-subtraction of the PSF leading to an upper-limit of the flux of C of $K_s=9.34$. The lines bisect the AB Dor A PSF and mark the location of AB Dor C.




Fig. 2.— The orbital solution of Nielsen et al. (2005) with our 2 new data points (#4 this work & #5 paper 1) added. Note that this orbit was mainly determined by the reflex motion of A from ~ 1 mas Hipparcos/VLBI astrometry of Guirado et al. (2006) (for clarity these Hipparcos/VLBI datapoints of A’s motion are not shown above – see Guirado et al. (2006) for details).




Fig. 3.— The separation of AB Dor C w.r.t. A as a function of time. The solid line is the orbital solution of Nielsen et al. (2005) with our new data points (the most recent two datapoints) added on top of the predicted orbit (this is not a new fit). The agreement is excellent with both the new Ks and IFS astrometric datapoints. Hence the mass of AB Dor C is confirmed to be within the range $0.090 \pm 0.005 M_{\odot}$.




Fig. 4.— The F_{λ} spectrum of AB Dor C (BLACK LINE – smoothed to template resolution) shown against a number of young (~ 5 Myr), low-surface gravity objects (colored lines; Gorlova et al. 2003). The features seem to be similar to the M6 template. The bottom curves are the residuals from the spectrum of AB Dor C and the various templates. Note the rather low S/N of these faint young templates and small errors in the extinction correction leads to some uncertainty in their continuum which makes a good spectral type determination difficult, hence more weight should be placed on the higher S/N field M dwarfs shown in figures 5 & 6.




Fig. 5.— The unsmoothed F_λ K spectrum of AB Dor C, this time plotted against high S/N field M dwarfs (~ 5 Gyr), with higher surface gravities (Cushing et al. 2005). A spectral type of $M5.5 \pm 1.0$ (1σ) seems most consistent with our spectrum since the gravity independent CO lines fit the M5 best but the continuum fits the M6 best. Note how the Na I lines are weaker than the M5 template but the CO fits very well, this is what we would expect for a young M5. The “emission” line at 2.28 microns is due to a poor telluric subtraction.




Fig. 6.— The unsmoothed F_λ H spectrum of AB Dor C, plotted against field M dwarfs (~ 5 Gyr), with higher surface gravities (Cushing et al. 2005). A spectral type of $M5.5 \pm 1.0$ (1σ) seems most consistent with our spectrum since the Mg & Al lines between 1.65 – $1.75 \mu\text{m}$ fit the M6 slightly better but the 1.57 – 1.8 micron continuum fits the M5 or the M6 equally well. The “emission” lines at 1.56 , 1.64 , and 1.74 microns are all due a poor telluric subtraction.




Fig. 7.— Comparison of our observations of AB Dor C (J&H from Close et al.(2005); K_s and T_{eff} from this work; all shown as solid squares; age 75 Myr) and those of Luhman & Potter (open triangles; plotted with their assumed older 120 Myr age) to the 0.095, 0.090, and 0.085 $M_{\odot}$ cooling curves from the DUSTY models of Chabrier et al. (2000). Note the good agreement with our new K_s and T_{eff} observations; while there is some overshoot in the J and H predicted fluxes for a 0.090 $M_{\odot}$ object.




Fig. 8.— HR diagram showing low-mass Pleiades objects from Martin et al. 2000 (open stars), other low-mass members of the Pleiades taken from the literature (open triangles), and AB Dor Ba/Bb (filled boxes). The dashed “vertical” lines are iso-mass contours for the DUSTY models (from left to right: 0.09, 0.07, 0.05, and 0.04 $M_{\odot}$), while the more “horizontal” lines are the DUSTY isochrones (top to bottom: 1, 10, 50, 100, 120, 500, 1000 Myr). Note that the DUSTY models predict a 75 Myr object of 0.09 $M_{\odot}$ should be similar in temperature and luminosity to that observed. From the location of AB Dor C on the HR diagram, one would derive a mass of $\sim 0.09 M_{\odot}$, a good estimate of the measured mass. As the temperature (dwarf T_{eff} scale of Leggett et al. 1996; Luhman 1999) and luminosities (BC_K of Allen et al. 2003) of the Pleiades objects in this plot were determined in the same manner used for AB Dor C, and these Pleiades points mostly fall along the appropriate 120 Myr DUSTY isochrone (dotted line), we are assured that our temperature scale and bolometric correction are reasonable. The points with error bars mark all known low-mass young objects with well determined dynamical masses, with the thick short diagonal lines representing the displacement from the measured luminosity and temperature to the values actually predicted by the DUSTY models. One can see that for AB Dor C and the older (300 Myr) Gl 569 B system (Zapatero Osorio et al. 2004), the offset between observation and the tracks are within the 1σ uncertainties (especially considering the uncertainty in the ages for these systems). There is somewhat less agreement for the secondary of the very young 2MASS 0535 eclipsing binary system in Orion (Stassun et al. 2006).